arXiv:1411.7429v1 [math.QA] 27 Nov 2014

Quantum dimensions and fusion rules for parafermion vertex operator algebras

Chongying Dong[a1] and Qing Wang[b2]
[a] Department of Mathematics, University of California, Santa Cruz, CA 95064
[b]School of Mathematical Sciences, Xiamen University, Xiamen 361005, China

Abstract

The quantum dimensions and the fusion rules for the parafermion vertex operator algebra associated to the irreducible highest weight module for the affine Kac-Moody algebra $A_1^{(1)}$ of level k are determined.

1 Introduction

Parafermion vertex operator algebra is the commutant of the Heisenberg vertex operator algebra in the affine vertex operator algebra. It comes from a special kind of coset construction [19]. Precisely speaking, let $L_{\widehat{\mathfrak{g}}}(k,0)$ be the level k integrable highest weight module with weight zero for affine Kac-Moody algebra $\widehat{\mathfrak{g}}$ associated to a finite dimensional simple Lie algebra $\mathfrak{g}$. Then $L_{\widehat{\mathfrak{g}}}(k,0)$ contains the Heisenberg vertex operator subalgebra generated by the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$. The commutant $K(\mathfrak{g},k)$ of the Heisenberg vertex operator subalgebra in $L_{\widehat{\mathfrak{g}}}(k,0)$ is the parafermion vertex operator algebra. The structure and representation theory of parafermion vertex operator algebras has been widely studied these years(see [11], [12], [14], [15], [3]). In particular, [14] and [15] show the role of parafermion vertex operator algebra $K(sl_2,k)$ in the parafermion vertex algebra $K(\mathfrak{g},k)$ is similar to the role of 3-dimensional simple Lie algebra sl_2 played in Kac-Moody Lie algebras, we denote $K(sl_2,k)$ by K_0 in this paper. Moreover, it was proved that K_0 coincides with a certain W-algebra in [11] and [12]. Later in [3], the C_2-cofiniteness of parafermion vertex operator algebra $K(\mathfrak{g},k)$ has been established by proving the C_2-cofiniteness of parafermion vertex operator algebra K_0, and irreducible modules for parafermion vertex operator algebra K_0 were also classified therein. In the recent paper [13], the rationality of parafermion vertex operator algebra $K(\mathfrak{g},k)$ was also obtained, moreover, the irreducible modules for $K(\mathfrak{g},k)$ were classified.

The notion of quantum dimensions of modules for vertex operator algebras was introduced in [6]. It was proved therein for rational and C_2-cofinite vertex operator algebras, quantum dimensions do exist. In this paper, we first determine the quantum dimensions for the parafermion vertex operator algebra K_0. Then by using the important formula obtained in [6] which shows that quantum dimensions are multiplicative under tensor product, we give the fusion rules for the parafermion vertex operator algebra K_0.

The paper is organized as follows. In Section 2, we recall some results about parafermion vertex operator algebra K_0. In Section 3, after reviewing the notion and properties of

[1]Supported by NSF grant DMS-1404741, NSA grant H98230-14-1-0118 and China NSF grant 11371261.

[2]Supported by China NSF grant (No.11371024), Natural Science Foundation of Fujian Province (No.2013J01018) and Fundamental Research Funds for the Central University (No.2013121001).

quantum dimensions of modules for vertex operator algebras, we give the quantum dimensions of parafermion vertex operator algebra K_0. In the final section, we obtained the fusion rules of parafermion vertex operator algebra K_0 by using the results of quantum dimensions of parafermion vertex operator algebra K_0.

2 Preliminary

In this section, we recall from [11], [12] and [3] some basic results on the parafermion vertex operator algebra associated to the irreducible highest weight module for the affine Kac-Moody algebra $A_1^{(1)}$ of level k with $k \geq 2$ being an integer. First we recall the notion of the parafermion vertex operator algebra.

We are working in the setting of [11]. Let $\{h, e, f\}$ be a standard Chevalley basis of sl_2 with brackets $[h, e] = 2e$, $[h, f] = -2f$, $[e, f] = h$. Let $\widehat{sl}_2 = sl_2 \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}C$ be the affine Lie algebra associated to sl_2. Let $k \geq 2$ be an integer and

$$V(k, 0) = V_{\widehat{sl}_2}(k, 0) = \mathrm{Ind}_{sl_2 \otimes \mathbb{C}[t] \oplus \mathbb{C}C}^{\widehat{sl}_2} \mathbb{C}$$

be an induced $\widehat{sl}_2$-module such that $sl_2 \otimes \mathbb{C}[t]$ acts as 0 and C acts as k on $\mathbf{1} = 1$. We denote by $a(n)$ the operator on $V(k, 0)$ corresponding to the action of $a \otimes t^n$. Then

$$[a(m), b(n)] = [a, b](m + n) + m\langle a, b\rangle \delta_{m+n,0} k \tag{1.1}$$

for $a, b \in sl_2$ and $m, n \in \mathbb{Z}$. It is well known that there is a vertex operator algebra structure on $V(k, 0)$ and it has a unique maximal ideal $\mathcal{J}$, which is generated by a weight $k + 1$ vector $e(-1)^{k+1}\mathbf{1}$ [21]. The quotient algebra $L(k, 0) = V(k, 0)/\mathcal{J}$ is the simple vertex operator algebra associated to an affine Lie algebra $\widehat{sl}_2$ of type $A_1^{(1)}$ with level k. The subspace $V_{\widehat{\mathfrak{h}}}(k, 0)$ of $V(k, 0)$ spanned by $h(-i_1) \cdots h(-i_p)\mathbf{1}$ for $i_1 \geq \cdots \geq i_p \geq 1$ and $p \geq 0$ is a vertex operator subalgebra of $V(k, 0)$ associated to the Heisenberg algebra. The parafermion vertex operator algebra K_0 is defined as the commutant of $V_{\widehat{\mathfrak{h}}}(k, 0)$ in $L(k, 0)$, that is,

$$K_0 = \{v \in L(k, 0) \,|\, h(m)v = 0 \text{ for } m \geq 0\}.$$

It was proved that K_0 is a simple vertex operator algebra and the irreducible K_0-modules $M^{i,j}$ for $0 \leq i \leq k, 0 \leq j \leq k - 1$ were constructed in [11]. Note that $K_0 = M^{0,0}$. It was also proved that $M^{i,j} \cong M^{k-i,k-i+j}$ as K_0-module in [11, Theorem 4.4] and moreover, Theorem 8.2 in [3] showed that the $\frac{k(k+1)}{2}$ irreducible K_0-modules $M^{i,j}$ for $0 \leq i \leq k, 0 \leq j \leq i - 1$ constructed in [11] form a complete set of isomorphism classes of irreducible K_0-module. Moreover, K_0 is C_2-cofinite [3] and rational [13].

Recall from [11] that $L = \mathbb{Z}\alpha_1 + \cdots + \mathbb{Z}\alpha_k$ with $\langle \alpha_p, \alpha_q\rangle = 2\delta_{pq}$. $V_L = M(1) \otimes \mathbb{C}[L]$ is the lattice vertex operator algebra associated with the lattice L. Let $\gamma = \alpha_1 + \cdots + \alpha_k$. Thus $\langle \gamma, \gamma\rangle = 2k$. It is well known that the vertex operator algebra associated with a positive definite even lattice is rational [5]. And any irreducible modules for the lattice vertex operator algebra $V_{\mathbb{Z}\gamma}$ is isomorphic to one of $V_{\mathbb{Z}\gamma+n\gamma/2k}$, $0 \leq n \leq 2k - 1$ [5]. Let $L(k, i)$ for $0 \leq i \leq k$ be the irreducible modules for the rational vertex operator algebra $L(k, 0)$. The following result was due to [11].

Lemma 2.1. $L(k,i) = \oplus_{j=0}^{k-1} V_{\mathbb{Z}\gamma+(i-2j)\gamma/2k} \otimes M^{i,j}$ *as* $V_{\mathbb{Z}\gamma}$*-modules.*

3 Quantum dimensions for irreducible K_0-modules

In this section, we first recall the notion and some basic facts about quantum dimension from [6]. Then we determine the quantum dimensions of the irreducible K_0-modules.

Let $(V, Y, 1, \omega)$ be a vertex operator algebra (see [18], [23]). We define weak module, module and admissible module following [7, 8]. Let $W\{z\}$ denote the space of W-valued formal series in arbitrary complex powers of z for a vector space W.

Definition 3.1. *A* weak V-module M *is a vector space with a linear map*

$$Y_M : V \to (EndM)\{z\}$$

$$v \mapsto Y_M(v,z) = \sum_{n\in\mathbb{Z}} v_n z^{-n-1} \quad (v_n \in EndM)$$

which satisfies the following conditions for $u, v \in V, w \in M$*:*

$$u_n w = 0 \text{ for } n \gg 0,$$

$$Y_M(\mathbf{1}, z) = Id_M,$$

$$z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right) Y_M(u,z_1) Y_M(v,z_2) - z_0^{-1}\delta\left(\frac{z_2 - z_1}{-z_0}\right) Y_M(v,z_2) Y_M(u,z_1)$$

$$= \delta\left(\frac{z_1 - z_0}{z_2}\right) Y_M(Y(u,z_0)v, z_2),$$

where $\delta(z) = \sum_{n\in\mathbb{Z}} z^n$.

Definition 3.2. *A* V-module *is a weak* V*-module* M *which carries a* $\mathbb{C}$*-grading* $M = \bigoplus_{\lambda\in\mathbb{C}} M_\lambda$*, where* $M_\lambda = \{w \in M | L(0)w = \lambda w\}$ *and* $L(0)$ *is one of the coefficient operators of* $Y(\omega, z) = \sum_{n\in\mathbb{Z}} L(n) z^{-n-2}$*. Moreover we require that* $\dim M_\lambda$ *is finite and for fixed* λ*,* $M_{\lambda+n} = 0$ *for all small enough integers* n*.*

Definition 3.3. *An* admissible V-module $M = \oplus_{n\in\mathbb{Z}_+} M(n)$ *is a* $\mathbb{Z}_+$*-graded weak module such that* $u_m M(n) \subset M(wtu - m - 1 + n)$ *for homogeneous* $u \in V$ *and* $m, n \in \mathbb{Z}$*.*

Definition 3.4. *A vertex operator algebra* V *is called* rational *if the admissible module category is semisimple.*

The following lemma about rational vertex operator algebras is well known [9].

Lemma 3.5. *If V is rational and M is an irreducible admissible V-module, then*

(1) M is a V-module and there exists a $\lambda \in \mathbb{C}$ such that $M = \oplus_{n\in\mathbb{Z}_+} M_{\lambda+n}$ where $M_\lambda \neq 0$. And λ is called the conformal weight of M;

(2) There are only finitely many irreducible admissible V-modules up to isomorphism.

Definition 3.6. *We say that a vertex operator algebra V is* C_2-cofinite *if $V/C_2(V)$ is finite dimensional, where $C_2(V) = \langle v_{-2}u | v, u \in V\rangle$.*

Let $M = \bigoplus_{\lambda\in\mathbb{C}} M_\lambda$ be a V-module. Set $M' = \bigoplus_{\lambda\in\mathbb{C}} M_\lambda^*$, the restricted dual of M where $M_\lambda^* = \mathrm{Hom}_{\mathbb{C}}(M_\lambda, \mathbb{C})$. It was proved in [16] that M' is naturally a V-module where the vertex operator $Y_{M'}(v, z)$ is defined for $v \in V$ via

$$\langle Y_{M'}(v, z)f, u\rangle = \langle f, Y_M(e^{zL(1)}(-z^{-2})^{L(0)}v, z^{-1})u\rangle,$$

where $\langle f, w\rangle = f(w)$ is the natural paring $M' \times M \to \mathbb{C}$. The V-module M' is called the *contragredient module* of M. A V-module M is called *self-dual* if M and M' are isomorphic V-modules. The following result was proved in [22].

Lemma 3.7. *Let V be a simple vertex operator algebra such that $L(1)V_{(1)} \neq V_{(0)}$. Then V is self-dual.*

Remark 3.8. *Note that the weight one subspace of K_0 is zero. By using lemma 3.7, parafermion vertex operator algebra K_0 is obviously self-dual.*

Now we recall from [16] the notions of intertwining operators and fusion rules.

Definition 3.9. *Let $(V,\ Y)$ be a vertex operator algebra and let $(W^1,\ Y^1)$, $(W^2,\ Y^2)$ and $(W^3,\ Y^3)$ be V-modules. An* intertwining operator *of type $\binom{W^3}{W^1\ W^2}$ is a linear map*

$$I(\cdot,\ z) :\ W^1 \to Hom(W^2,\ W^3)\{z\}$$

$$u \to I(u,\ z) = \sum_{n\in\mathbb{Q}} u_n z^{-n-1}$$

satisfying:

(1) for any $u \in W^1$ and $v \in W^2$, $u_n v = 0$ for n sufficiently large;

(2) $I(L(-1)v,\ z) = \frac{d}{dz} I(v,\ z)$;

(3) (Jacobi identity) for any $u \in V,\ v \in W^1$

$$z_0^{-1}\delta\left(\frac{z_1 - z_2}{z_0}\right) Y^3(u,\ z_1)I(v,\ z_2) - z_0^{-1}\delta\left(\frac{-z_2 + z_1}{z_0}\right) I(v,\ z_2)Y^2(u,\ z_1)$$
$$= z_2^{-1}\left(\frac{z_1 - z_0}{z_2}\right) I(Y^1(u,\ z_0)v,\ z_2).$$

The space of all intertwining operators of type $\binom{W^3}{W^1\ W^2}$ is denoted by

$$I_V\binom{W^3}{W^1\ W^2}.$$

Let $N_{W^1,\ W^2}^{W^3} = \dim I_V \begin{pmatrix} W^3 \\ W^1 \ \ W^2 \end{pmatrix}$. *These integers* $N_{W^1,\ W^2}^{W^3}$ *are usually called the* fusion rules.

Definition 3.10. *Let* V *be a vertex operator algebra, and* W^1, W^2 *be two* V*-modules. A module* (W, I)*, where* $I \in I_V \begin{pmatrix} W \\ W^1 \ \ W^2 \end{pmatrix}$*, is called a* tensor product *(or fusion product) of* W^1 *and* W^2 *if for any* V*-module* M *and* $\mathcal{Y} \in I_V \begin{pmatrix} M \\ W^1 \ \ W^2 \end{pmatrix}$*, there is a unique* V*-module homomorphism* $f : W \to M$*, such that* $\mathcal{Y} = f \circ I$*. As usual, we denote* (W, I) *by* $W^1 \boxtimes_V W^2$.

Remark 3.11. *It is well known that if* V *is rational, then for any two irreducible* V*-modules* W^1 *and* W^2*, the fusion product* $W^1 \boxtimes_V W^2$ *exists and*

$$W^1 \boxtimes_V W^2 = \sum_W N_{W^1,\ W^2}^W W,$$

where W *runs over the set of equivalence classes of irreducible* V*-modules.*

Now we recall some notions about quantum dimensions.

Definition 3.12. *Let* $M = \oplus_{n \in \mathbb{Z}_+} M_{\lambda+n}$ *be a* V*-module, the* formal character *of* M *is defined as*

$$ch_q M = tr_M q^{L(0)-c/24} = q^{\lambda - c/24} \sum_{n \in \mathbb{Z}_+} (\dim M_{\lambda+n})\, q^n,$$

where c *is the central charge of the vertex operator algebra* V *and* λ *is the conformal weight of* M.

It is proved [24, 10] that $\mathrm{ch}_q M$ converges to a holomorphic function in the domain $|q| < 1$. We denote the holomorphic function $\mathrm{ch}_q M$ by $Z_M(\tau)$. Here and below, τ is in the upper half plane $\mathbb{H}$ and $q = e^{2\pi i \tau}$.

Let $M^0, \cdots, M^d$ be the inequivalent irreducible V-modules with corresponding conformal weights λ_i and $M^0 \cong V$.

From the Remark 2.13 of [6], we have

Definition 3.13.

$$Z_{M^i}\left(-\frac{1}{\tau}\right) = \sum_{j=0}^{d} S_{i,j} Z_{M^j}(\tau).$$

The matrix $S = (S_{i,j})_{i,j=0}^d$ *is called an* S-matrix.

The following definition of quantum dimension was introduced in [6].

Definition 3.14. *Let V be a vertex operator algebra and M a V-module such that $Z_V(\tau)$ and $Z_M(\tau)$ exist. The quantum dimension of M over V is defined as*

$$qdim_V M = \lim_{y\to 0} \frac{Z_M(iy)}{Z_V(iy)},$$

where y is real and positive.

The following result was obtained in [6, Lemma 4.2].

Lemma 3.15. *Let V be a simple, rational and C_2-cofinite vertex operator algebra of CFT type with $V \cong V'$. Let M^i for $0 \le i \le d$ be the inequivalent irreducible V-modules with corresponding conformal weights λ_i and $M^0 \cong V$. Assume $\lambda_0 = 0$ and $\lambda_i > 0$ $\forall i \neq 0$. Then $qdim_V M^i = \frac{S_{i,0}}{S_{0,0}}$.*

From now on, we assume V is a rational, C_2-cofinite vertex operator algebra of CFT type with $V \cong V'$. Let $M^0 \cong V$, M^1, $\cdots$, M^d denote all inequivalent irreducible V-modules. Moreover, we assume the conformal weights λ_i of M^i are positive for all $i > 0$. From Remark 3.8 and statement in Section 2, the parafermion vertex operator algebra K_0 satisfies all the assumptions.

The following result shows that the quantum dimensions are multiplicative under tensor product [6] .

Proposition 3.16. *Let V and M_i for $0 \le i \le d$ be as in Lemma 3.15. Then*

$$qdim_V\left(M^i \boxtimes M^j\right) = qdim_V M^i \cdot qdim_V M^j$$

for i, $j = 0, \cdots, d$.

Before giving the main result of this section, we recall the following character of irreducible K_0-modules $M^{i,j}$ which is given in [4, 20]:

$$\mathrm{ch}M^{i,j} = \eta(\tau) c^i_{i-2j}(\tau)$$

by [20, (3.34)]. Note that k, l and m in [20] are k, i and $i-2j$, respectively in our notation.

Theorem 3.17. *The quantum dimensions for all irreducible K_0-modules $M^{m,n}$ are*

$$qdim_{K_0} M^{m,n} = \frac{\sin\frac{\pi(m+1)}{k+2}}{\sin\frac{\pi}{k+2}}$$

for $0 \le m \le k$, $0 \le n \le k-1$, where $M^{m,n}$ are the irreducible modules of K_0 constructed in [11].

Proof. Let $M^{m,n}(\tau)$ denote the character of $M^{m,n}$ for $0 \le m \le k$, $0 \le n \le k-1$. The S-modular transformation of characters has the following form [20], [21]:

$$M^{m,n}(-\frac{1}{\tau}) = \sum_{m',n'} S^{m',n'}_{m,n} M^{m',n'}(\tau),$$

where $S^{m^{'},n^{'}}_{m,n} = (k(k+2))^{-\frac{1}{2}} \exp \frac{i\pi(m-2n)(m^{'}-2n^{'})}{k} \sin \frac{\pi(m+1)(m^{'}+1)}{k+2}$. From [11] and [3], we see that K_0 has $\frac{k(k+2)}{2}$ irreducible modules $M^{m,n}$ with the conformal weights

$$\lambda_{m,n} = \frac{1}{2k(k+2)}(k(m-2n) - (m-2n)^2 + 2kn(m-n+1))$$

for $0 \leq m \leq k,\ 0 \leq n \leq m-1$. It is easy to check that $\lambda_{k,0} = 0$ and $\lambda_{m,n} > 0$ for $(m,n) \neq (k,0)$. Thus by using Lemma 3.15, we have

$$\mathrm{qdim}_{K_0} M^{m,n} = \frac{S^{0,0}_{m,n}}{S^{0,0}_{0,0}} = \frac{\sin \frac{\pi(m+1)}{k+2}}{\sin \frac{\pi}{k+2}}.$$

□

4 Fusion rule for irreducible K_0-modules

In this section, we give the fusion rules for irreducible K_0-modules. First we fix some notations. Let W^1, W^2, W^3 be irreducible K_0-modules. In the following, we use $I \begin{pmatrix} W^3 \\ W^1\, W^2 \end{pmatrix}$ to denote the space $I_{K_0} \begin{pmatrix} W^3 \\ W^1\, W^2 \end{pmatrix}$ of all intertwining operators of type $\begin{pmatrix} W^3 \\ W^1\, W^2 \end{pmatrix}$, and use $W^1 \boxtimes W^2$ to denote the fusion product $W^1 \boxtimes_{K_0} W^2$ for simplicity.

We recall the fusion rules for affine vertex operator algebra of type $A_1^{(1)}$ [**?**] for later use.

Lemma 4.1.

$$L(k,i) \boxtimes_{L(k,0)} L(k,j) = \sum_c L(k,c),$$

where $|i-j| \leq c \leq i+j,\ i+j+c \in 2\mathbb{Z},\ i+j+c \leq 2k$.

Theorem 4.2. *The fusion rule for the irreducible modules of Parafermion vertex operator algebra K_0 is as follows:*

$$M^{i,i^{'}} \boxtimes M^{j,j^{'}} = \sum_c M^{c,\overline{\frac{1}{2}(2i^{'}-i+2j^{'}-j+c)}}, \tag{4.1}$$

where $\overline{a}$ means the residue of the integer a modulo k, $0 \leq i,j \leq k, 0 \leq i^{'}, j^{'} \leq k-1$, $|i-j| \leq c \leq i+j,\ i+j+c \in 2\mathbb{Z},\ i+j+c \leq 2k$. Moreover, with fixed $i, i^{'}, j, j^{'}$, $M^{c,\overline{\frac{1}{2}(2i^{'}-i+2j^{'}-j+c)}}$ for $|i-j| \leq c \leq i+j,\ i+j+c \in 2\mathbb{Z},\ i+j+c \leq 2k$ are inequivalent irreducible modules.

Proof. We take $V = L(k,0), U = V_{\mathbb{Z}\gamma} \otimes K_0$ in Proposition 2.9 of [1], from the Lemma 2.1, we see that

$$\dim I_V \begin{pmatrix} L(k,l) \\ L(k,i)\, L(k,j) \end{pmatrix} \leq \dim I_U \begin{pmatrix} L(k,l) \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \otimes M^{i,i^{'}}\ V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \otimes M^{j,j^{'}} \end{pmatrix}$$

for $0 \leq i, j, l \leq k, 0 \leq i^{'}, j^{'}, l^{'} \leq k-1$. Note that $L(k,l) = \oplus_{j=0}^{k-1} V_{\mathbb{Z}\gamma+(l-2j)\gamma/2k} \otimes M^{l,j}$, thus we have

$$\dim I_U \begin{pmatrix} L(k,l) \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \otimes M^{i,i^{'}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \otimes M^{j,j^{'}} \end{pmatrix}$$

$$= \sum_{l^{'}=0}^{k-1} \dim I_U \begin{pmatrix} V_{\mathbb{Z}\gamma+\frac{(l-2l^{'})\gamma}{2k}} \otimes M^{l,l^{'}} \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \otimes M^{i,i^{'}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \otimes M^{j,j^{'}} \end{pmatrix}.$$

Then by using Theorem 2.10 of [1], we have

$$\dim I_U \begin{pmatrix} V_{\mathbb{Z}\gamma+\frac{(l-2l^{'})\gamma}{2k}} \otimes M^{l,l^{'}} \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \otimes M^{i,i^{'}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \otimes M^{j,j^{'}} \end{pmatrix}$$

$$= \dim I_{V_{\mathbb{Z}_\gamma}} \begin{pmatrix} V_{\mathbb{Z}\gamma+\frac{(l-2l^{'})\gamma}{2k}} \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \end{pmatrix} \cdot \dim I_{K_0} \begin{pmatrix} M^{l,l^{'}} \\ M^{i,i^{'}} \; M^{j,j^{'}} \end{pmatrix}.$$

Recall that the fusion rule for lattice vertex operator algebras is:

$$V_{\mathbb{Z}\gamma+\lambda} \boxtimes_{V_{\mathbb{Z}_\gamma}} V_{\mathbb{Z}\gamma+\mu} = V_{\mathbb{Z}\gamma+\lambda+\mu}$$

for $\lambda, \mu \in (\mathbb{Z}\gamma)^\circ$, where $(\mathbb{Z}\gamma)^\circ$ is the dual lattice of $\mathbb{Z}\gamma$, this together with the fusion rule for affine vertex operator algebra given in Lemma 4.1, we see that if $l^{'} \neq \frac{l-i+2i^{'}-j+2j^{'}}{2}$, then

$$\dim I_U \begin{pmatrix} V_{\mathbb{Z}\gamma+\frac{(l-2l^{'})\gamma}{2k}} \otimes M^{l,l^{'}} \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \otimes M^{i,i^{'}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \otimes M^{j,j^{'}} \end{pmatrix}$$

$$= \dim I_{V_{\mathbb{Z}_\gamma}} \begin{pmatrix} V_{\mathbb{Z}\gamma+\frac{(l-2l^{'})\gamma}{2k}} \\ V_{\mathbb{Z}\gamma+\frac{(i-2i^{'})\gamma}{2k}} \; V_{\mathbb{Z}\gamma+\frac{(j-2j^{'})\gamma}{2k}} \end{pmatrix} \cdot \dim I_{K_0} \begin{pmatrix} M^{l,l^{'}} \\ M^{i,i^{'}} \; M^{j,j^{'}} \end{pmatrix} = 0.$$

And we also have

$$\dim I_{K_0} \begin{pmatrix} M^{c,\overline{\frac{1}{2}(2i^{'}-i+2j^{'}-j+c)}} \\ M^{i,i^{'}} \; M^{j,j^{'}} \end{pmatrix} \geq 1.$$

In the following, we use the quantum dimension of the irreducible modules over K_0 to prove that

$$\dim I_{K_0} \begin{pmatrix} M^{c,\overline{\frac{1}{2}(2i^{'}-i+2j^{'}-j+c)}} \\ M^{i,i^{'}} \; M^{j,j^{'}} \end{pmatrix} = 1.$$

Since from Proposition 3.16, we know

$$\mathrm{qdim}_{K_0} \left(M^{i,i^{'}} \boxtimes M^{j,j^{'}} \right) = \mathrm{qdim}_{K_0} M^{i,i^{'}} \cdot \mathrm{qdim}_{K_0} M^{j,j^{'}},$$

and we also have

$$M^{i,i'} \boxtimes M^{j,j'} = \sum_{(0,0)\leq(l,l')\leq(k,l-1)} N^{(l,l')}_{(i,i'),(j,j')} M^{l,l'}.$$

Since Theorem 3.17 shows that

$$\mathrm{qdim}_{K_0} M^{i,i'} = \frac{\sin\frac{\pi(i+1)}{k+2}}{\sin\frac{\pi}{k+2}}.$$

Thus we only need to prove the following identity holds:

$$\frac{\sin\frac{\pi(i+1)}{k+2}\cdot\sin\frac{\pi(j+1)}{k+2}}{\sin\frac{\pi}{k+2}\cdot\sin\frac{\pi}{k+2}} = \sum_c \frac{\sin\frac{\pi(c+1)}{k+2}}{\sin\frac{\pi}{k+2}}$$

for $0 \leq i,j \leq k, 0 \leq i',j' \leq k-1$, $|i-j| \leq c \leq i+j,\ i+j+c \in 2\mathbb{Z},\ i+j+c \leq 2k$. This identity is equivalent to the following identity:

$$\cos\frac{\pi(i+j+2)}{k+2} - \cos\frac{\pi(i-j)}{k+2} = \sum_c (\cos\frac{\pi(c+2)}{k+2} - \cos\frac{\pi c}{k+2}). \tag{4.2}$$

We note that if $i+j<k$, then $c_{\min} = i-j$, $c_{\max} = i+j$, thus (4.2) holds. If $i+j>k$, $c_{\min} = i-j$, $c_{\max} = i+j-2n$ for some n satisfying that $i+j-2n+i+j = 2k$, that is, $c_{\max} = 2k-i-j$, thus (4.2) also holds.

Now we prove that the modules $M^{c,\overline{\frac{1}{2}(2i'-i+2j'-j+c)}}$ appeared in the sum (4.1) are not isomorphic to each other. If $M^{c,\overline{\frac{1}{2}(2i'-i+2j'-j+c)}} \cong M^{c',\overline{\frac{1}{2}(2i'-i+2j'-j+c')}}$ for some c,c' satisfying $|i-j| \leq c \leq i+j,\ i+j+c \in 2\mathbb{Z},\ i+j+c \leq 2k$. Since we know that $M^{i,j} \cong M^{k-i,k-i+j}$ as K_0-module for $0 \leq i \leq k, 0 \leq j \leq k-1$ and the $\frac{k(k+1)}{2}$ irreducible K_0-modules $M^{i,j}$ for $0 \leq i \leq k, 0 \leq j \leq i-1$ exhaust all the isomorphism classes of irreducible K_0-modules, it follows that $c' = k-c$ and

$$\overline{k-c+\frac{1}{2}(2i'-i+2j'-j+k-c)} = \overline{\frac{1}{2}(2i'-i+2j'-j+k-c)},$$

it is impossible by a direct calculation. This proves the assertion.

□

References

[1] T. Abe, C. Dong and H. Li, Fusion rules for the vertex operator $M(1)^+$ and V_L^+, *Comm. Math. Phys.*. **253** (2005) 171-219.

[2] T. Arakawa, Representation theory of W-algebras, *Invent. Math.* **169** (2007), 219–320.

[3] T. Arakawa, C.H. Lam and H. Yamada, Zhu's algebra, C_2-cofiniteness of parafermion vertex operator algebras, *Adv. Math.* **264** (2014), 261–295.

[4] A. Cappelli, L. S. Georgiev and I. T. Todorov, Parafermion Hall states from coset projections of abelian conformal theories, *Nucl. Phys.* **B599** (2001), 499–530.

[5] C. Dong, Vertex algebras associated with even lattices, *J. Algebra* **160** (1993), 245–265.

[6] C. Dong, X. Jiao and F. Xu, Quantum dimensions and Quantum Galois theory, *Trans. AMS.* **365** (2013), 6441-6469.

[7] C. Dong, H. Li and G. Mason, Simple currents and extensions of vertex operator algebras. *Comm. Math. Phys.* **180** (1996) 671-707.

[8] C. Dong, H. Li and G. Mason, Regularity of rational vertex operator algebras, *Adv. Math.* **132** (1997) 148-166.

[9] C. Dong, H. Li and G. Mason, Twisted representations of vertex operator algebras, *Math. Ann.* **310** (1998) 571-600.

[10] C. Dong, H. Li and G. Mason, Modular-Invariance of Trace Functions in Orbifold Theory and Generalized Moonshine, *Comm. Math. Phys.* **214** (2000) 1-56.

[11] C. Dong, C.H. Lam and H. Yamada, W-algebras related to parafermion algebras, *J. Algebra* **322** (2009), 2366-2403.

[12] C. Dong, C.H. Lam, Q. Wang and H. Yamada, The structure of parafermion vertex operator algebras, *J. Algebra* **323** (2010), 371-381.

[13] C. Dong and L. Ren, Representations of the parafermion vertex operator algebras, arXiv:1411.6085.

[14] C. Dong and Q. Wang, The structure of parafermion vertex operator algebras: general case, *Comm. Math. Phys.* **299** (2010), 783-792.

[15] C. Dong and Q. Wang, On C_2-cofiniteness of parafermion vertex operator algebras, *J. Algebra* **328** (2011), 420-431.

[16] I. B. Frenkel, Y. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, *Memoirs American Math. Soc.* **104**, 1993.

[17] E. Frenkel, V. Kac and M. Wakimoto, Characters and fusion rules for W-algebras via quantized Drinfeld-Sokolov reduction, *Comm. Math. Phys.* **147** (1992), 295-328.

[18] I. B. Frenkel, J. Lepowsky and A. Meurman, Vertex operator algebras and the monster, *Pure and Applied Math.* Vol. 134, Academic Press, Massachusetts, 1988.

[19] P. Goddard, A. Kent and D. Olive, Unitary representations of the Virasoro and super Virasoro algebras, *Comm. Math. Phys.* **103** (1986), 105–119.

[20] D. Gepner and Z. Qiu, Modular invariant partition functions for parafermionic field theories, *Nucl. Phys.* **B285** (1987), 423–453.

[21] V. G. Kac, *Infinite-dimensional Lie Algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.

[22] H. Li, Symmetric invariant bilinear forms on vertex operator algebras, *J. pure Appl. Alg.* **96** (1994) 279-297.

[23] J. Lepowsky and H. Li, *Introduction to Vertex Operator Algebras and Their Representations*, Progress in Math., Vol. 227, Birkhäuser, Boston, 2004.

[24] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996) 237-302.